CHILD BIRTH

| NAME: | DATE: |
|-------|-------|
| | |

| EXAMINATION STEPS AND PROCEDURE | PC | P0 | COMMENTS |
|---|-------|--------|------------------|
| Introduce self, reassure patient and request permission to treat | | 1 | |
| 2. Ask patient. "Are you under a doctor's case'?" | | 1 | |
| 3. Ask patient, "Does your doctor expect any problems. with the delivery?" | | 10 | |
| 4. Ask patient. "Is this your first vaginal delivery?" | | 10 | |
| 5 Ask patient, "When did the contractions begin?" | | 10 | |
| 6 Ask patient, "How far apart are the contractions?" | | 10 | |
| 7. Ask patient, "Do you feel a need to bear down or move your bowels?" | | 10 | |
| 8. Reassure patient of need to examine for crowning: | | 1 | |
| 9Observe for crowning | | 20 | |
| BABY'S HEAD SEEN.CROWNING AND IS NOW A FACE DOWN (NORMAL DELIVERY) | ADVAN | CING (| OUT, SLOWLY WITH |
| 10. Open OB kit | | 1 | |
| 11 Attempt to cleanse area and drape | | 1 | |
| 12. Put gloves on | | 1 | |
| 13. Use sterile technique | | 10 | |
| 14. With one hand, apply gentle pressure to baby's head to . prevent sudden expulsion . | | 20 | |
| 15. As the head emerges, using other hand; apply gentle downward pressure to the perineum directly below the vaginal opening to prevent tearing | | 10 | |

| EXAMINATION STEPS AND PROCEDURE | PC | P0 | COMMENTS | | | | |
|--|---|----|----------|--|--|--|--|
| Report to scorer presence or absence of cord around baby's neck when baby's head is out | | 10 | | | | | |
| 17. If present. loosen cord from around baby's neck | | 10 | | | | | |
| 18. Slip cord over baby's head | | 10 | | | | | |
| 19 Clear airway by sucti9ning baby's mouth with bulb syringe:A. Squeeze air out of bulb syringe before entering orifice | | 10 | | | | | |
| B Suction nose | | 10 | | | | | |
| C. Suction mouth | | 10 | | | | | |
| 20. Apply gentle downward pressure to head to release upper shoulder | | 1 | | | | | |
| 21 Apply gentle upward -pressure to head to release lower shoulder | | 1 | | | | | |
| BODY DELIVERING QUICKLY NOW | | | | | | | |
| 22 Hold baby securely with firm but gentle grip | | 20 | | | | | |
| 23. Clear airway by suctioning: baby's mouth with bulb syringe: A Squeeze air out of bulb syringe before entering orifice | | 10 | | | | | |
| B. Suction nose | | 10 | | | | | |
| C. Suction mouth | | 10 | | | | | |
| SCORER STATES, "BABY IS OUT, NOT BREATH | SCORER STATES, "BABY IS OUT, NOT BREATHING BUT HAS PULSE" | | | | | | |
| 24. Stimulate baby by: | | | | | | | |
| A. Rubbing back briskly or B Flicking bottom of baby's foot | | 30 | | | | | |

| EXAMINATION STEPS AND PROCEDURE | PC | P0 | COMMENTS | | | |
|---|----|----|----------|--|--|--|
| SCORER STATES, "BABY IS NOW CRYING' | | 20 | | | | |
| 25. Wrap baby in clean blanket for warmth | | 1 | | | | |
| 26. Place baby on mother's abdomen or between legs | | 10 | | | | |
| 27. Double damp cord: A. First clamp should be six to eight inches from baby | | 10 | | | | |
| B. Second clamp should be two to four inches from first clamp towards mother | | 10 | | | | |
| SCORER STATES, "PLACENTA HAS DELIVERED." | | | | | | |
| 28. Place placenta in plastic bag in preparation for transport | | 1 | | | | |
| 29. Externally massage uterus | | 20 | | | | |
| 30. Skill completed within 10 minutes | | 10 | | | | |

Scorer's Signature